

# Leading Axion-Photon Sensitivity with NuSTAR Observations of M82 and M87

Orion Ning  
University of California, Berkeley  
Lawrence Berkeley National Laboratory

*with Benjamin Safdi*  
arXiv:2404.14476

## Key Highlights:

1. New axion-photon constraints using NuSTAR observations of M82/M87 galaxies
2. Use of full galaxies as a probe of axion physics

# A Tale of Two Galaxies



M82



M87



# A Tale of Two Galaxies



M82

- Starburst Galaxy
- Indications of Strong B-fields



M87

# A Tale of Two Galaxies



M82


- Starburst Galaxy
- Indications of Strong B-fields



M87

- Massive Elliptical
- Extended Virgo Cluster B-fields


# Searching for Axions with NuSTAR:



# Searching for Axions with NuSTAR:

Axions produced via Primakoff in stellar interiors


$$g_{a\gamma\gamma}$$



# Searching for Axions with NuSTAR:

Axions produced via Primakoff in stellar interiors

$$g_{a\gamma\gamma}$$




Convert to hard X-rays in galactic/cluster magnetic fields



# Searching for Axions with NuSTAR:

Axions produced via Primakoff in stellar interiors

$$g_{a\gamma\gamma}$$



Convert to hard X-rays in galactic/cluster magnetic fields

Observed by the NuSTAR telescope

# Axion Signal Model Ingredients

$$\frac{dF}{dE}(E) = P_{a \rightarrow \gamma}(E) \frac{1}{4\pi d^2} \frac{dL_a(E)}{dE}$$

Axion-Photon Conversion Probability

Distance

Total Axion Luminosity from Stars

# Signal Model Ingredients (1/2)

Axion Luminosity from Stellar Populations of M82/M87

# Signal Model Ingredients (1/2)

## Axion Luminosity from Stellar Populations of M82/M87

Primakoff Process  $\frac{dL_a(E)}{dE}$



# Signal Model Ingredients (1/2)

## Axion Luminosity from Stellar Populations of M82/M87

Primakoff Process



Stellar Profiles (MESA)

$$\frac{dL_a(E)}{dE}$$

- Temperature
- Density
- Abundances

# Signal Model Ingredients (1/2)

## Axion Luminosity from Stellar Populations of M82/M87

Primakoff Process



Stellar Profiles (MESA)



Stellar Population  
Models (Obs.)

$$\frac{dL_a(E)}{dE}$$

- Temperature
- Density
- Abundances
  
- Metallicity
- SFH
- IMF
- # of Stars

# Signal Model Ingredients (1/2)

## Axion Luminosity from Stellar Populations of M82/M87

Primakoff Process



Stellar Profiles (MESA)



Stellar Population Models (Obs.)




Total Axion Luminosity Spectra

$$\frac{dL_a(E)}{dE}$$

- Temperature
- Density
- Abundances
  
- Metallicity
- SFH
- IMF
- # of Stars

- Dominated by Supergiants and O-Types, generally
- M82: starburst = excellent for axions!



# Signal Model Ingredients (2/2)

Conversion Probability for Axion-Photon Conversion



# Signal Model Ingredients (2/2)


## Conversion Probability for Axion-Photon Conversion

Axion Mass

$m_a$

+

Magnetic Fields



+

Free-electron Densities

$n_e$


||

Conversion  
Probabilities

$P_{a \rightarrow \gamma}$


# Signal Model Ingredients (2/2)

## Conversion Probability for Axion-Photon Conversion




# Signal Model Ingredients (2/2)

## Conversion Probability for Axion-Photon Conversion



The variability in galaxy candidates and magnetic fields = largest source of uncertainty in our analysis

# No evidence for axions from NuSTAR = Upper limits on coupling





Previous Limits: C. O'Hare



In Progress: Can M82 give strong limits for other couplings?


# In Progress: Can M82 give strong limits for other couplings?

- Tentative strong limits for axion-electron and axion-nucleon couplings from bremsstrahlung production
- Highlights NS population in latter case




# Conclusion and Outlook

- Leading constraints on the axion-photon coupling from X-ray observations of all stars in M82/M87




# Conclusion and Outlook

- Leading constraints on the axion-photon coupling from X-ray observations of all stars in M82/M87
- Galaxies as a probe of axion physics




# Conclusion and Outlook

- Leading constraints on the axion-photon coupling from X-ray observations of all stars in M82/M87
- Galaxies as a probe of axion physics
- Magnetic fields dominant source of uncertainty



# Conclusion and Outlook

- Leading constraints on the axion-photon coupling from X-ray observations of all stars in M82/M87
- Galaxies as a probe of axion physics
- Magnetic fields dominant source of uncertainty
- Can extend to other galaxies, clusters, and axion-electron and axion-nucleon couplings





Thanks for listening!





# Appendix





# Signal Model + NuSTAR Data Constrains Axion-Photon Coupling




# Systematics







# Stellar Population Modeling



# Axion Luminosity for M87



# Conversion Probabilities



# Spatial Maps

