


Observations of the IC 443 region with HAWC

Hugo Ayala
Penn State University

TeV Particle Astrophysics, Chicago, 2024



Supernova remnants as Cosmic-ray accelerators



- Supernova Remnants (SNRs) are the aftermaths of the catastrophic explosions of massive stars or white dwarfs after accreting enough mass
- They are capable of accelerating cosmic rays (CRs)
- Candidates to be PeVatrons, i.e. CRs at PeV energies.


IC 443 Nebula



- One of the most studied SNRs. Observed for the first time in 1892.
- Located at ~ 1.8 kpc (Ambrocio-Cruz et al. 2017)
- Detected in radio, IR, optical, x-rays and gamma rays:
 - Radio highlights regions of synchrotron radiation (Castelletti et al. 2011, and others).
 - Optical helps delineate the boundaries and the region of the interaction of the shock wave and the ISM (Li et al. 2022, and others)
 - X-rays reveal the hot gas heated by the shock wave. Also found the progenitor of the SNR, a pulsar named CXOU J061705.3+222127 (Swartz et al. 2015)
 - Gamma rays alludes to particle acceleration and type of particles (Ackermann et al. 2013, and others)



High Altitude Water Cherenkov Observatory



Energy range: ~ 300 GeV — 100 TeV
Angular resolution: $\sim 0.2^\circ$
Field of View: ~ 2 sr
>95% Uptime

Main array completed March 2015
Outriggers deployed 2018


4100 m elevation

100,000 m²

22,000 m²



HAWC NIM paper 2023

HAWC observations of the IC 443 region



HAWC observations of the IC 443 region


- Contours from Lee, Jae-Joon, et al. 2008.






- Angular resolution of HAWC is ~ 0.2 deg above 10 TeV.

Analysis of the region


- Use likelihood method (3ML, Vianello et al. 2015; Abeysekara et al. 2022)
 - Find morphology and spectrum that best describes the data:
 - Iterative process similar to Fermi-LAT catalog construction
 - Best model: a point source close to IC 443 and an extended source (Gaussian shape).
 - Both prefer a power-law spectrum.
 - Includes Galactic Diffuse Emission from Hermes model (Dundovic et al. 2021)



Analysis - Model and residual maps




IC 443 - VHE





- Fermi confirmed hadronic acceleration after observing pionic gamma rays (2013)
- HAWC observations increase the energy range of the observed gamma rays
 - PL with index of -3.14 and normalization of $(5.9 \pm 1.3) \times 10^{-13}$ $[\text{TeV cm}^2 \text{s}]^{-1}$ at 1 TeV.
- Non-thermal emission model prefers a broken power law function for the CR spectra.
- LHAASO flux from catalog. Their search found one extended source. A detailed analysis is needed for proper comparison.
- Not enough evidence to tell if it is a PeVatron

IC 443 - MW picture




- Non-thermal emission model
- IR data is thermal emission so it's not part of the final model, but used as a photon seed for inverse Compton.
- X-ray observations are from the PWN so are not considered in the Fit

IC443 - MW picture



- In this lepto-hadronic model, IC and Pion decay both contribute to the VHE emission above 10 TeV.

Extended source - a new TeV Halo?




- The extended emission is around PSR B0611+22 (or J0614+2229)
- Old pulsar with 89kyr of age
- Located at 3.55kpc away
- $\dot{E} = 6.24 \times 10^{34} \text{ erg s}^{-1}$
- $P_0 = 0.3349 \text{ s}$


VHE Halos in a nutshell

- “Inverse Compton” halos: gamma rays emitted by inverse Compton scattering of electrons and positrons accelerated in the pulsar wind nebulae

- A couple of models:




Sudoh et al. 2019



Giacinti et al. 2020

- Test evolution of particle acceleration and escape from PWN and pulsars
- Study propagation of electrons and positrons in the ISM


Extended source - a new TeV Halo?




- Simple inverse Compton model
- Use the fitted electron spectrum to find energy density of electrons
- Smaller than the ISM -> Evidence that this is a electron (TeV) halo.
- Current estimates with HAWC data put this value between 10^{-3} and $10^{-5} \text{ eV cm}^{-3}$

Extended source - CR illumination of Gas?

- Another hypothesis, accelerated CR interacting with gas.
- Diffuse gamma-ray emission from sea of CRs interacting with the gas included in the model
- Fresh CRs accelerated in the region could produce this emission. Calculations in progress.



HI (HI4PI survey)



H2 (Dame CO survey)

Caveat: integrated over full velocity range

Summary

- HAWC Observations of IC 443 up to 40 TeV.
 - Consistent with other VHE observations. Data can be described by a lepto-hadronic model
- An extended source is observed in the region. Emission could come from
 - Halo around pulsar B0611+22
 - CR illumination of gas



Thank you




Image credit: Giuseppe Donatiello

Back-up

Numbers

Table 1. Results of the analysis in the region of IC 443 using the f_{hit} scheme.

| Source | K [TeV ⁻¹ cm ⁻² s ⁻¹] | Index | R.A. [deg] | Decl. [deg] | σ [deg] |
|--------|--|------------------------------------|-------------------------------------|-------------------------------------|------------------------------------|
| IC 443 | $(5.9 \pm 1.3^{+0.21}_{-0.86}) \times 10^{-14}$ | $-3.14 \pm 0.18^{+0.08}_{-0.09}$ | $94.42^{+0.07+0.009}_{-0.05-0.008}$ | $22.35^{+0.06+0.035}_{-0.07-0.003}$ | — |
| E.S. | $(3.18^{+1.37+0.19}_{-0.92-0.37}) \times 10^{-13}$ | $-2.49 \pm 0.08^{+0.009}_{-0.028}$ | $93.67 \pm 0.19^{+0.016}_{-0.004}$ | $22.22 \pm 0.20^{+0.015}_{-0.007}$ | $1.05^{+0.21+0.004}_{-0.18-0.013}$ |

Naima Models

- IC 443 - Hadronic Model:
 - SBPL: $\log(N)=47.34+0.09-0.06$; $\text{idx1} = 2.36\pm0.04$; $\text{idx2} = 3.04\pm0.05$; $E_{\text{break}} = 100\pm30$ GeV
 - ECPL: $\log(N)=49.58\pm0.03$; $\text{idx1} = 2.34\pm0.03$; $E_{\text{cutoff}} = 71\pm9$ GeV
- IC 443 - Lepto Hadronic Model:
 - SBPL proton: $\log(N)=47.11+5.6-0.1$; $\text{idx1} = 2.35\pm0.08$; $\text{idx2} = 3.15\pm0.2$; $E_{\text{break}} = 174\pm30$ GeV; $N_{\text{H}} = 188\pm12$ cm²
 - SBPL electron: $\log(N)=48.6\pm0.2$; $\text{idx1} = 1.68\pm0.04$; $\text{idx2} = 3.61\pm0.2$; $E_{\text{break}} = 50\pm15$ GeV; $B=10.4\pm1.0$ uG
- Halo - IC model:
 - SPL = $\log(N)=47.7\pm0.3$; $\text{idx}=3.42\pm0.18$
 - ECPL = $\log(N)=47.7\pm0.5$; $\text{idx}=2.87\pm0.73$; $E_{\text{cutoff}} = 66\pm20$ TeV

Importance of Halos

- Test evolution of particle acceleration and escape from PWN and pulsars
- Study propagation of electrons and positrons in the ISM

Halo Comparisons

